

HAMILTONIANS REPRESENTING EQUATIONS OF MOTION WITH DAMPING DUE TO FRICTION

STEPHEN MONTGOMERY-SMITH

ABSTRACT. Suppose that $H(q, p)$ is a Hamiltonian on a manifold M , and $\tilde{L}(q, \dot{q})$, the Rayleigh dissipation function, satisfies the same hypotheses as a Lagrangian on the manifold M . We provide a Hamiltonian framework that gives the equation

$$\dot{q} = \frac{\partial H}{\partial p}(q, p), \quad \dot{p} = -\frac{\partial H}{\partial q}(q, p) - \frac{\partial \tilde{L}}{\partial \dot{q}}(q, \dot{q})$$

The method is to embed M into a larger framework where the motion drives a wave equation on the negative half line, where the energy in the wave represents heat being carried away from the motion. We obtain a version of Nöther's Theorem that is valid for dissipative systems. We also show that this framework fits the widely held view of how Hamiltonian dynamics can lead to the "arrow of time."

1. INTRODUCTION

The purpose of this document is to provide a Hamiltonian framework which gives rise to equations of motion that include friction or damping effects. The author has found a number of other works [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12], but none of them seem to solve the problem in the manner presented here. The methods of this paper allow for extremely general formulas for the damping term, but the method is also artificial. Nevertheless we are able to extract various principles such as Nöther's Theorem.

2. NOTATION

We represent the position of a particle by q , which might be in a finite or infinite dimensional manifold M . (Everything in this document is formal, and there is no attempt at rigor.) As usual we have a Lagrangian $L(q, \dot{q})$ which is strictly convex in the second coordinate. The equations of motion are the solution to the variational equation $\delta\mathcal{S} = 0$, where the action \mathcal{S} is

$$\mathcal{S} = \int_{T_0}^{T_1} L(q, \dot{q}) dt; \tag{2.1}$$

2000 *Mathematics Subject Classification.* 70H25.

Key words and phrases. Hamiltonian; Lagrangian; Rayleigh dissipation function; friction; Nöther's Theorem.

©2014 Texas State University - San Marcos.

Submitted January 23, 2014. Published April 2, 2014.

that is,

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right) - \frac{\partial L}{\partial q} = 0 \quad (2.2)$$

As usual, we create the Hamiltonian

$$H(q, p) = p \cdot \dot{q} - L(q, \dot{q}) \quad (2.3)$$

where p satisfies the equation

$$p = \frac{\partial L}{\partial \dot{q}} \quad (2.4)$$

It can then be shown that

$$\dot{q} = \frac{\partial H}{\partial p} \quad (2.5)$$

$$\frac{\partial L}{\partial q} = - \frac{\partial H}{\partial q} \quad (2.6)$$

where in the latter formula one has to be aware that the first partial derivative is keeping \dot{q} constant, and the second partial derivative is keeping p constant. The equation of motion becomes

$$\dot{p} = - \frac{\partial H}{\partial q} \quad (2.7)$$

The usual example is

$$L(q, \dot{q}) = \frac{1}{2} \dot{q} \cdot A \cdot \dot{q} - V(q) \quad (2.8)$$

where $A = A(q)$ is positive definite symmetric. Then

$$p = A \cdot \dot{q} \quad (2.9)$$

$$H(q, p) = \frac{1}{2} p \cdot A^{-1} \cdot p + V(q) \quad (2.10)$$

and the equation of motion is

$$\frac{d}{dt} (A \cdot \dot{q}) - \frac{1}{2} \dot{q} \cdot \frac{\partial A}{\partial q} \cdot \dot{q} + \frac{\partial V}{\partial q} = 0 \quad (2.11)$$

The book [1] gives many more details.

3. STRIVING TO UNDERSTAND FRICTION

Here we describe a widely held view of how friction arises from reversible Hamiltonian dynamics. Our concrete equations of motion obtained in Sections 4 and 5 should conform to the framework described here.

Suppose the manifold M is a submanifold of $M_0 \times M_1$, where M_0 represents the macroscopic motion, and M_1 represents the microscopic motion. The position then splits as $q = (q_0, q_1)$, and the momentum splits as $p = p_0 + p_1$ in the obvious manner. Because M_1 represents microscopic motion, it makes sense that M_1 should have much higher dimension than M_0 , indeed we will often suppose that M_1 is infinite dimensional. For example, maybe the microscopic motion is a large number of high frequency harmonic oscillators.

We assume that the Hamiltonian splits as

$$H(q, p) = H_0(q_0, p_0) + H_1(q_0, q_1, p_1) \quad (3.1)$$

Here H_0 represents the macroscopic motion without friction. The second part, H_1 , is the energy of the microscopic motion plus coupling between the microscopic and macroscopic parts. We assume that all the interaction between the microscopic and macroscopic parts is via position and not momentum.

Because H_1 does not depend upon p_0 , from equation (2.5) we obtain

$$\dot{q}_0 = \frac{\partial H_0}{\partial p_0} \quad (3.2)$$

$$\dot{q}_1 = \frac{\partial H_1}{\partial p_1} \quad (3.3)$$

and the Lagrangian splits as

$$L(q, \dot{q}) = L_0(q_0, \dot{q}_0) + L_1(q_0, q_1, \dot{q}_1) \quad (3.4)$$

where

$$L_0(q_0, \dot{q}_0) = p_0 \cdot \dot{q}_0 - H_0(q_0, p_0) \quad (3.5)$$

$$L_1(q_0, q_1, \dot{q}_1) = p_1 \cdot \dot{q}_1 - H_1(q_0, q_1, p_0) \quad (3.6)$$

and equations (2.4) and (2.6) become

$$p_0 = \frac{\partial L_0}{\partial \dot{q}_0} \quad (3.7)$$

$$p_1 = \frac{\partial L_1}{\partial \dot{q}_1} \quad (3.8)$$

$$\frac{\partial L_0}{\partial q} = -\frac{\partial H_0}{\partial q} \quad (3.9)$$

$$\frac{\partial L_1}{\partial q} = -\frac{\partial H_1}{\partial q} \quad (3.10)$$

(The only mildly non-obvious statements are equation (3.9) which follows from $\partial L_0/\partial q_0 = -\partial H_0/\partial q_0$ and $\partial L_0/\partial q_1 = \partial H_0/\partial q_1 = 0$, and equation (3.10) which follows by subtraction.)

The equations of motion now become

$$\dot{p}_0 = -\frac{\partial H_0}{\partial q_0} - \frac{\partial H_1}{\partial q_0} \quad (3.11)$$

$$\dot{p}_1 = -\frac{\partial H_1}{\partial q_1} \quad (3.12)$$

Now, if the microscopic motion starts at rest, then we *hope* that the second law of thermodynamics should be obeyed, and as a function of time, H_0 should be non-increasing, and H_1 should be non-decreasing. Thus the hope is that the motion described by q_1 somehow provides a memory of what has happened to q_0 , so that we end up with

$$\frac{\partial H_1}{\partial q_0} = \Phi(q_0, \dot{q}_0) \quad (3.13)$$

for some function Φ . Then H_0 satisfies

$$\begin{aligned} \dot{H}_0 &= \dot{q}_0 \cdot \frac{\partial H_0}{\partial q_0} + \dot{p}_0 \cdot \frac{\partial H_0}{\partial p_0} \\ &= \dot{q}_0 \cdot (-\dot{p}_0 - \Phi(q_0, \dot{q}_0)) + \dot{p}_0 \cdot \dot{q}_0 = -\dot{q}_0 \cdot \Phi(q_0, \dot{q}_0) \end{aligned} \quad (3.14)$$

If Φ satisfies a property like

$$y \cdot \Phi(x, y) \geq 0 \quad (3.15)$$

then we that H_0 is non-increasing in time.

The case we can handle with our approach is to consider

$$\Phi(x, y) = \frac{\partial \tilde{L}}{\partial y}(x, y) \quad (3.16)$$

where $y \mapsto \tilde{L}(x, y)$ is a strictly convex. Note that property (3.15) holds if $\tilde{L}(x, y)$ attains its minimum at $y = 0$, because

$$\begin{aligned} y \cdot \frac{\partial \tilde{L}}{\partial y}(x, y) &= y \cdot \frac{\partial \tilde{L}}{\partial y}(x, 0) + \int_0^1 \frac{\partial}{\partial \lambda} \left(y \cdot \frac{\partial \tilde{L}}{\partial y}(x, \lambda y) \right) d\lambda \\ &= \int_0^1 y \cdot \frac{\partial^2 \tilde{L}}{\partial y^2}(x, \lambda y) \cdot y d\lambda \geq 0 \end{aligned} \quad (3.17)$$

since the Hessian of $\tilde{L}(x, \cdot)$ is positive definite.

The initial values of q_1 and p_1 should be very important, perhaps representing that the parts in M_1 are at rest, or are oscillating in such a manner that the oscillations can only increase. This is because Hamiltonian equations are reversible in time. But dissipative systems should only be solvable forwards in time. This is sometimes known as the ‘‘arrow of time.’’ For example, we know that if a wine glass is thrown to the ground, then it shatters. The reverse process, where the wine glass comes back together, is possible in theory if the initial positions and momenta of the molecules and atoms are given in a very precise manner (and we neglect the effects of quantum physics). But finding this initial data should be very difficult.

4. FRICTION VIA THE WAVE EQUATION: PROOF OF CONCEPT

We will show how to artificially create damping via the wave equation. As a proof of concept, let us explain how to create a Hamiltonian such that

$$\dot{q}_0 = \frac{\partial H_0}{\partial p_0} \quad (4.1)$$

$$\dot{p}_0 = -\frac{\partial H_0}{\partial q_0} - B \cdot \dot{q}_0 \quad (4.2)$$

for some positive definite matrix B .

An example of an infinite dimensional Hamiltonian is the wave equation on $C^1(\mathbb{R}, M)$, whose Hamiltonian is given by

$$\int_{-\infty}^{\infty} \frac{1}{2} p(s) \cdot B^{-1} \cdot p(s) + \frac{1}{2} q'(s) \cdot B \cdot q'(s) ds \quad (4.3)$$

Here q' denotes $\partial q / \partial s$. Now the equation of motion coming from this Hamiltonian is

$$B\ddot{q} = Bq'' \quad (4.4)$$

which by d'Alembert's principle has a general solution

$$q(s, t) = \phi_1(s + t) + \phi_2(s - t) \quad (4.5)$$

This is a good example of a Hamiltonian for which Poincaré's recurrence theorem does not apply. And furthermore, the equations of motion are a point of variation of the Lagrangian, but are clearly not a minimum of the Lagrangian:

$$\int_{-\infty}^{\infty} \frac{1}{2} \dot{q}(s) \cdot B \cdot \dot{q}(s) - \frac{1}{2} q'(s) \cdot B \cdot q'(s) ds \quad (4.6)$$

The way we simulate friction is to consider the macroscopic motion as driving a wave equation on the half line. Let $M_1 = C^1((-\infty, 0], M_0)$, that is, the set of functions $q : (-\infty, 0] \rightarrow M_0$ such that q' is bounded, $q'(s) \rightarrow q'(0)$ as $s \rightarrow 0^-$, where $q'(0)$ denotes the left derivative of $q(s)$ at $s = 0$, and $q'(s) \rightarrow 0$ as $s \rightarrow -\infty$.

Then let

$$M = \{(q_0, q_1) \in M_0 \times M_1 : q_1(0) = q_0\} \quad (4.7)$$

with the Hamiltonian

$$H(q, p) = H_0(q_0, p_0) + \int_{-\infty}^0 \frac{1}{2} p_1(s) \cdot B^{-1} \cdot p_1(s) + \frac{1}{2} q_1'(s) \cdot B \cdot q_1'(s) ds \quad (4.8)$$

We see that the equation of motion is

$$\dot{q}_0 = \frac{\partial H_0}{\partial p_0} \quad (4.9)$$

$$\dot{p}_0 = -\frac{\partial H_0}{\partial q_0} - B \cdot q_1'(0) \quad (4.10)$$

$$B\ddot{q}_1 = Bq_1'' \quad (4.11)$$

For example, to obtain equations (4.9) and (4.11), see that for any infinitesimal perturbation $\delta q = (\delta q_0, \delta q_1)$ of $q = (q_0, q_1)$, noting that $\delta q_0 = \delta q_1(0)$ and $\delta q_1'(s) \rightarrow 0$ as $s \rightarrow -\infty$, we have

$$\begin{aligned} \delta q \cdot \frac{\partial H}{\partial q} &= \delta q_0 \cdot \frac{\partial H_0}{\partial q_0}(q_0, p_0) + \int_{-\infty}^0 \delta q_1'(s) \cdot B \cdot q_1'(s) ds \\ &= \delta q_0 \cdot \frac{\partial H_0}{\partial q_0}(q_0, p_0) + \left[\delta q_1(s) \cdot B \cdot q_1'(s) \right]_{-\infty}^0 - \int_{-\infty}^0 \delta q_1(s) \cdot B \cdot q_1''(s) ds \\ &= \delta q_0 \cdot \frac{\partial H_0}{\partial q_0}(q_0, p_0) + \delta q_0 \cdot B \cdot q_1'(0) - \int_{-\infty}^0 \delta q_1(s) \cdot B \cdot q_1''(s) ds \end{aligned}$$

Next we impose initial conditions on (q_1, p_1)

$$q_1(s, T_0) = q_0(T_0), \quad p_1(s, T_0) = 0 \quad (s \leq 0) \quad (4.12)$$

that is, the microscopic part of the motion is initially at rest. Then it can be seen that the solution to the wave equation is given by (4.5) with $\phi_1(t) = q_0(t)$ and $\phi_2 = 0$, that is

$$q_1(s, t) = \begin{cases} q_1(s+t-T_0, T_0) = q_0(T_0) & \text{if } s+t \leq T_0 \\ q_0(s+t) & \text{if } s+t \geq T_0 \end{cases} \quad (4.13)$$

In particular, we see that $q_1'(0, t) = \dot{q}_1(0, t) = \dot{q}_0(t)$, and hence we obtain equations (4.1) and (4.2).

Thus it is seen that the irreversibility of equations of motion with damping comes naturally from the special nature of the initial conditions. (And indeed any initial condition satisfying $q_1'(s, T_0) = \dot{q}_1(s, T_1)$ and $q_1(0, T_0) = q_0(T_0)$ will work just as well.)

5. THE GENERAL EQUATION OF MOTION WITH DAMPING

Suppose that \tilde{H} , and its corresponding Lagrangian \tilde{L} , have the same domains as H_0 and L_0 . Here \tilde{L} is also called the Rayleigh dissipation function.

Consider the Hamiltonian on $M = \{(q_0, q_1) \in M_0 \times M_1 : q_1(0) = q_0\}$ given by

$$H(q, p) = H_0(q_0, p_0) + \int_{-\infty}^0 \tilde{H}(q_1(s), p_1(s)) + \tilde{L}(q_1(s), q_1'(s)) ds \quad (5.1)$$

Note that the corresponding Lagrangian is

$$L(q, \dot{q}) = L_0(q_0, \dot{q}_0) + \int_{-\infty}^0 \tilde{L}(q_1(s), \dot{q}_1(s)) - \tilde{L}(q_1(s), q_1'(s)) ds \quad (5.2)$$

Theorem 5.1. *The equations of motion for the Hamiltonian given by (5.1) with initial conditions (4.12) imply*

$$\dot{q}_0 = \frac{\partial H_0}{\partial p_0} \quad (5.3)$$

$$\dot{p}_0 = -\frac{\partial H_0}{\partial q_0}(q_0, p_0) - \frac{\partial \tilde{L}}{\partial \dot{q}_0}(q_0, \dot{q}_0) \quad (5.4)$$

Proof. First, looking at the equations of motion for (q_1, p_1) we obtain

$$\dot{p}_1 = -\frac{\partial \tilde{H}}{\partial q_1}(q_1, p_1) - \frac{\partial \tilde{L}}{\partial q_1}(q_1, q_1') + \frac{\partial}{\partial s} \left(\frac{\partial \tilde{L}}{\partial q_1'}(q_1, q_1') \right) \quad (5.5)$$

$$\dot{q}_1 = \frac{\partial \tilde{H}}{\partial p_1}(q_1, p_1) \quad (5.6)$$

where the last term of equation (5.5) comes by integrating by parts, just as in the previous section.

From the definition of \tilde{L} , we see that equation (5.6) is equivalent to

$$p_1 = \frac{\partial \tilde{L}}{\partial \dot{q}_1}(q_1, \dot{q}_1) \quad (5.7)$$

and substituting into equation (5.5) we obtain

$$\frac{\partial}{\partial t} \left(\frac{\partial \tilde{L}}{\partial \dot{q}_1}(q_1, \dot{q}_1) \right) - \frac{\partial \tilde{L}}{\partial q_1}(q_1, \dot{q}_1) = \frac{\partial}{\partial s} \left(\frac{\partial \tilde{L}}{\partial q_1'}(q_1, q_1') \right) - \frac{\partial \tilde{L}}{\partial q_1}(q_1, q_1') \quad (5.8)$$

This is a wave equation, and since we have initial conditions (4.12), the solution (4.13) is valid. (The general solution is not so obvious: if $\tilde{L}(x, y) = \tilde{L}(x, -y)$, then $\phi_1(s+t)$ and $\phi_2(s-t)$ are both solutions to this equation. However it is not obvious to me how these two solutions should be combined, since this wave equation is non-linear.)

Next, the equations of motion for (q_0, p_0) become

$$\dot{p}_0 = -\frac{\partial H_0}{\partial q_0}(q_0, p_0) - \frac{\partial \tilde{L}}{\partial q_1'(0)}(q_1(0), q_1'(0)) \quad (5.9)$$

$$\dot{q}_0 = \frac{\partial H_0}{\partial p_0}(q_0, p_0) \quad (5.10)$$

where the last term of equation (5.9) from the cross term from integrating by parts. From equation (4.13) we see that $q_1' = \dot{q}_1$. Substituting this into equation (5.9), and noting that $q_1(0) = q_0$, we obtain the desired result. \square

Finally, it is worth noting that under the initial conditions (4.12) the Hamiltonian and Lagrangian evaluate to

$$H(q(t), p(t)) = H_0(q_0(t), p_0(t)) + \int_{T_0}^t \left(\frac{\partial \tilde{L}}{\partial \dot{q}_0}(q_0(\tau), \dot{q}_0(\tau)) \right) \cdot \dot{q}_0(\tau) d\tau \tag{5.11}$$

$$L(q(t), \dot{q}(t)) = L_0(q_0(t), \dot{q}_0(t)) \tag{5.12}$$

In the case that $\tilde{L}(x, y) = \frac{1}{2}y \cdot B \cdot y$, the Hamiltonian becomes

$$H(q(t), p(t)) = H_0(q_0(t), p_0(t)) + \int_{T_0}^t \dot{q}_0(\tau) \cdot B \cdot \dot{q}_0(\tau) d\tau \tag{5.13}$$

6. THE ARROW OF TIME

Now that we have an explicit Hamiltonian that generates the equations of motion with friction, we should examine whether it satisfies the comments made at the end of Section 3 on the ‘‘arrow if time.’’

One way to reverse the effect of friction would be to first solve the equation backwards in time, from $t = T_1$ back to $t = T_0$, using initial conditions $q_1(s, T_1) = q_0(T_1)$, $p_1(s, T_1) = 0$ for $(s \leq 0)$. Solve to find $q_1(s, T_0)$, noting that it will satisfy $q'_1(s, T_0) = -\dot{q}_1(s, T_0)$. Then this is the initial data that is required to make the ‘wine glass come back together.’ Obviously not any initial data satisfying $q'_1(s, T_0) = -\dot{q}_1(s, T_0)$ will work, because the wave has to feed back into the equations of motion for q_0 exactly the energy it needs to satisfy the reverse effect of friction.

It would be interesting to see to what extent this equation is sensitive to the initial data, perhaps illustrating that it is very hard to get it exactly right to get the ‘wine glass to reconstruct.’ The idea would be to analyze the equations of motion for the full system on M , but either restricting to solutions in which $\dot{q}_1 = q'_1$ (in which case the solutions should be stable), or to the case in which $\dot{q}_1 = -q'_1$ (in which case the solutions should be unstable — this latter case requires $\tilde{L}(x, y) = \tilde{L}(x, -y)$, otherwise the formulation of the problem would be harder). Almost surely one will find that in the latter case, the equation is linearly unstable, corresponding to the fact that the Hessian of $\tilde{L}(x, \cdot)$ is positive definite.

7. NÖTHER’S THEOREM

By applying Nöther’s Theorem to our Lagrangian, we can easily obtain a version of Nöther’s Theorem that applies to dissipative systems. Suppose there is a flow on M_0 that preserves both L_0 and \tilde{L} , let us denote the parameter driving the flow by λ . Then the following quantity is conserved.

$$\pi(t) = \frac{\partial L_1}{\partial \dot{q}_0}(q_0(t), \dot{q}_0(t)) \cdot \frac{\partial q_0(t)}{\partial \lambda} \Big|_{\lambda=0} + \int_{T_0}^t \frac{\partial \tilde{L}}{\partial \dot{q}_0}(q_0(\tau), \dot{q}_0(\tau)) \cdot \frac{\partial q_0(\tau)}{\partial \lambda} \Big|_{\lambda=0} d\tau \tag{7.1}$$

Let us illustrate with a couple of examples of computing the momentum. Suppose that q_0 is n vectors in \mathbb{R}^m : $q_0 = (q_{01}, q_{02}, \dots, q_{0n})$. Let μ_x be the momentum in the direction x

$$\mu(t) = \sum_i m_i \dot{q}_{0i} \cdot x \tag{7.2}$$

which corresponds to the action

$$\lambda \mapsto (q_0 \mapsto (q_{0i} + \lambda x)_{i=1}^n) \tag{7.3}$$

Suppose that

$$L_0(q_0, \dot{q}_0) = \sum_i \frac{1}{2} m_i |\dot{q}_{0i}|^2 + \sum_i \sum_j \frac{1}{2} v_{ij} (|q_{0i} - q_{0j}|) \quad (7.4)$$

$$\tilde{L}(q_0, \dot{q}_0) = \eta \sum_i m_i \frac{1}{2} |\dot{q}_{0i}|^2 \quad (7.5)$$

where $v_{ij} = v_{ji}$. This gives the equation of motion

$$m_i \ddot{q}_{0i} = - \sum_j \frac{(q_{0i} - q_{0j})}{|q_{0i} - q_{0j}|} v'_{ij} (|q_{0i} - q_{0j}|) - \eta m_i \dot{q}_{0i} \quad (7.6)$$

Both of these Lagrangians are invariant under the action (7.3). We see that

$$\pi(t) = \mu_x(t) + \eta \int_{T_0}^t \mu_x(\tau) d\tau \quad (7.7)$$

is conserved; that is,

$$\dot{\mu}_x + \eta \mu_x = 0 \Rightarrow \mu_x(t) = e^{-\eta t} \mu_x(0) \quad (7.8)$$

Now suppose we change equation (7.5) to

$$\tilde{L}(q_0, \dot{q}_0) = \sum_i \sum_j \frac{1}{2} \eta_{ij} \left(\frac{\partial}{\partial t} |q_{0i} - q_{0j}| \right) \quad (7.9)$$

where $\eta_{ij} = \eta_{ji}$, that is, all the damping effects are internal and depend completely on how the distances between the particles change. The equation of motion is now

$$m_i \ddot{q}_{0i} = - \sum_j \frac{(q_{0i} - q_{0j})}{|q_{0i} - q_{0j}|} v'_{ij} (|q_{0i} - q_{0j}|) - \sum_j \frac{(q_{0i} - q_{0j})}{|q_{0i} - q_{0j}|} \eta'_{ij} \left(\frac{\partial}{\partial t} |q_{0i} - q_{0j}| \right) \quad (7.10)$$

This Lagrangian is also invariant under the action (7.3). We see that

$$\frac{\partial \tilde{L}}{\partial \dot{q}_0} (q_0(t), \dot{q}_0(t)) \cdot \frac{\partial q_0(t)}{\partial \lambda} \Big|_{\lambda=0} = \sum_i \sum_j \frac{(q_{0i} - q_{0j})}{|q_{0i} - q_{0j}|} \eta'_{ij} \left(\frac{\partial}{\partial t} |q_{0i} - q_{0j}| \right) = 0 \quad (7.11)$$

Therefore, $\pi = \mu_x$, and thus the momentum μ_x is conserved.

Let us try a similar argument with angular momentum. Let α_A be the angular momentum about the anti-symmetric matrix A

$$\alpha(t) = \sum_i m_i \dot{q}_{0i} \cdot A \cdot q_{0i} \quad (7.12)$$

which corresponds to the action

$$\lambda \mapsto (q_0 \mapsto (e^{\lambda A} \cdot q_{0i})_{i=1}^n) \quad (7.13)$$

Since $e^{\lambda A}$ is an orthogonal matrix, all of the above Lagrangians are invariant under the action (7.3). For the case (7.5)

$$\pi(t) = \alpha_A(t) + \eta \int_{T_0}^t \alpha_A(\tau) d\tau \quad (7.14)$$

is conserved; that is,

$$\dot{\alpha}_A + \eta \alpha_A = 0 \Rightarrow \alpha_A(t) = e^{-\eta t} \alpha_A(0) \quad (7.15)$$

For the case (7.9), we obtain

$$\frac{\partial \tilde{L}}{\partial \dot{q}_0}(q_0(t), \dot{q}_0(t)) \cdot \frac{\partial q_0(t)}{\partial \lambda} \Big|_{\lambda=0} = \sum_i \sum_j \frac{(q_{0i} - q_{0j})}{|q_{0i} - q_{0j}|} \eta'_{ij} \left(\frac{\partial}{\partial t} |q_{0i} - q_{0j}| \right) \cdot A \cdot q_{0i} = 0 \quad (7.16)$$

and thus similarly α_A is conserved.

(We should add that technically Lagrangian (7.9) does not quite fit into our framework, because it is not strictly convex in \dot{q}_0 . But this technicality is just that, and easily overcome. One can either redo all the work, or add $\epsilon|\dot{q}_0|^2$ to this Lagrangian and take the limit as $\epsilon \rightarrow 0$.)

8. AN IDENTITY

In this section we present an identity which we believe might be useful for creating conserved quantities. Suppose that equation (5.8) is satisfied. Let

$$E = \tilde{H}(p_1, q_1) - \tilde{L}(q_1, \dot{q}_1) \\ F = \dot{q}_1 \cdot \frac{\partial \tilde{L}}{\partial \dot{q}_1}(q_1, \dot{q}_1)$$

Then

$$\frac{\partial E}{\partial t} = \frac{\partial F}{\partial s} \quad (8.1)$$

In the case that $\tilde{L}(x, y) = y \cdot B(x) \cdot y$, so that $\tilde{L}(q_1, \dot{q}_1) = \tilde{H}(q_1, p_1)$, there is a symmetry in the formulas so that we also have

$$\frac{\partial E}{\partial s} = \frac{\partial F}{\partial t} \quad (8.2)$$

and hence both E and F satisfy the standard wave equation:

$$\frac{\partial^2 E}{\partial t^2} = \frac{\partial^2 E}{\partial s^2}, \quad \frac{\partial^2 F}{\partial t^2} = \frac{\partial^2 F}{\partial s^2} \quad (8.3)$$

REFERENCES

- [1] Arnold, V.I.; *Mathematical Methods of Classical Mechanics*, 2nd Ed, Springer-Verlag, 1989.
- [2] Bruneau, L.; De Bièvre, S.; *A Hamiltonian Model for Linear Friction in a Homogeneous Medium*, Communications in Mathematical Physics, September 2002, Volume 229, Issue 3, pp 511-542.
- [3] Chandrasekar, V. K.; Senthilvelan, M.; Lakshmanan, M.; *On the Lagrangian and Hamiltonian description of the damped linear harmonic oscillator*, J. Math. Phys. 48, 032701 (2007).
- [4] Fröhlich, J.; Gang, Z.; Soffer, A.; *Some Hamiltonian models of friction*, J. Math. Phys. 52, 083508 (2011).
- [5] Herreral, L.; Núñez, L.; Patiño, A.; Rago, H.; *A variational principle and the classical and quantum mechanics of the damped harmonic oscillator*, American Journal of Physics, March 1986, Volume 54, Issue 3, pp. 273.
- [6] Koopman, J.; Jeltsema, S.; Verhaegen, M.; *Port-Hamiltonian description and analysis of the LuGre friction model*, Simulation Modelling Practice and Theory - Simul Model Pract Theory, vol. 19, no. 3, pp. 959-968, 2011.
- [7] Rabei, E. M.; Alhalholy, T. S.; Taani, A. A.; *On Hamiltonian Formulation of Non-Conservative Systems*, Turk. J. Phys 28 (2004), 213-221.
- [8] Riewe, F.; *Nonconservative Lagrangian and Hamiltonian mechanics*, Phys. Rev. E 53, 1890-1899 (1996).
- [9] Riewe, F.; *Mechanics with fractional derivatives* Physical Review E, 55, (1997), 3581.
- [10] Smith, C. E.; *Lagrangians and Hamiltonians with friction*, J. Phys.: Conf. Ser. 237 (2010).

- [11] Yu, L. H.; Sun, C. P.; *Evolution of the wave function in a dissipative system*, Physical Review A, 49, (1994), 592.
- [12] Zhmoginov A.I.; Dodin, I.Y.; Fisch, N. J.; *A Hamiltonian model of dissipative wave-particle interactions and the negative-mass effect*, Physics Letters A 375 (2011) 1236-1241.

STEPHEN MONTGOMERY-SMITH
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA
E-mail address: stephen@missouri.edu